

Determine Bleed Severity

- Determining bleed severity is a key step in making treatment decisions.
- Bleeds can be classified into major and non-major based on several clinical factors.
- **If one or more of the following factors apply, the bleed should be considered major.**

For additional information, visit
www.anticoagulationtoolkit.org

Bleeding in critical site (examples below)	Hemodynamic instability (examples below)	Overt bleeding with either:
<ul style="list-style-type: none"> • Central nervous system bleeds (intracranial, spinal, intraocular) • Pericardial tamponade • Airway, including posterior epistaxis • Hemothorax 	<ul style="list-style-type: none"> • Intra-abdominal • Retroperitoneal • Intra-articular • Intra-muscular 	<ul style="list-style-type: none"> • Elevated heart rate • Decrease in SBP >40 mm Hg • Mean arterial pressure (intra-arterial) <65 mm Hg
	<ul style="list-style-type: none"> • SBP <90 mm Hg • Orthostatic blood pressure changes • Urine output <0.5 mL/kg/hr 	<ul style="list-style-type: none"> • Hemoglobin drop of ≥2 g/dL or • Administration of ≥2 U of packed RBCs

Assess for Clinically Relevant Drug Levels

- If last dose taken at least 24 hr ago in patients with normal renal function, drug levels probably not clinically relevant.¹
- If patient taking dabigatran, a TT can be used to rule out clinically relevant drug levels. Specialized tests can quantify drug levels.
- If apixaban, edoxaban, rivaroxaban, or betrixaban, anti-Xa is the preferred test and can be used to rule out relevant drug levels or quantify levels.
- **Don't wait for results before administering reversal agents in life-threatening bleeds!**

	Specialized Test	Drug Level Interpretation	General Test	Drug Level Interpretation
Dabigatran	dTT, ECT, ECA	Normal: not clinically relevant Results correlate with drug level	TT	Normal: not clinically relevant Prolonged: may/may not be clinically relevant
			aPTT	Normal: likely indicates lower drug level but can't exclude drug presence Prolonged: clinically relevant
Apixaban Betrixaban Edoxaban Rivaroxaban	Anti-Xa	Absent activity: not clinically relevant Results correlate with drug level (if calibrated for specific DOAC)	PT	Normal: does not exclude clinically relevant levels Prolonged: clinically relevant levels

Anti-Xa= anti-factor Xa; aPTT= activated partial thromboplastin time; dTT= dilute thrombin time; ECA= ecarin chromogenic assay; ECT= ecarin clotting time; PT= prothrombin time; TT= thrombin time

Manage Bleeding

All bleeds	Major Bleeds		Minor Bleeds	
	Critical site or life threatening	NOT critical site or life threatening	More serious minor bleeds†	Less serious minor bleeds
<ul style="list-style-type: none"> • Provide local therapy/ manual compression • Assess for and manage comorbidities contributing to the bleed* 	<ul style="list-style-type: none"> • Stop DOAC • Provide supportive care <ul style="list-style-type: none"> • Secure airway and large-bore IV access • Aggressive volume resuscitation (NS or LR) • Correct hypothermia and acidosis • Early involvement of other services (eg. surgery) • RBC transfusions to achieve Hgb ≥7 g/dL (≥8 g/dL if pt has CAD) • Platelet transfusion to achieve counts >50 x 10⁹/L • Cryoprecipitate transfusion to maintain fibrinogen >100 mg/dL • Stop any antiplatelets • Consider surgical/procedural management • Administer reversal agent 	<ul style="list-style-type: none"> • Stop DOAC • Provide supportive care <ul style="list-style-type: none"> • Secure airway and large-bore IV access • Aggressive volume resuscitation (NS or LR) • Correct hypothermia and acidosis • Early involvement of other services (eg. surgery) • RBC transfusions to achieve Hgb ≥7 g/dL (≥8 g/dL if pt has CAD) • Platelet transfusion to achieve counts >50 x 10⁹/L • Cryoprecipitate transfusion to maintain fibrinogen >100 mg/dL • Stop any antiplatelets • Consider surgical/procedural management • Administer reversal agent if above not effective 	<ul style="list-style-type: none"> • Stop DOAC • Provide supportive care • Stop any antiplatelets • Consider surgical/procedural management 	<ul style="list-style-type: none"> • Consider continuing DOAC if appropriate indication • Assess risk/benefits of stopping any antiplatelets • Verify that DOAC dosing is correct and patient taking as directed

*eg. renal dysfunction, liver disease, thrombocytopenia; † Patient requires hospitalization, transfusion, or procedural intervention

DOAC Reversal

Dabigatran	Apixaban, Betrixaban, Edoxaban, Rivaroxaban
<ul style="list-style-type: none"> • Administer 5 g idarucizumab IV (two separate 2.5 g/50 mL vials) <ul style="list-style-type: none"> • If bleeding persists and there is laboratory evidence of persistent dabigatran effect after 12-24 hours, a second dose may be reasonable. • If idarucizumab not available, administer aPCC (4F-PCC if aPCC not available) 50 units/kg IV (refer to package insert for max units)⁴ • Activated charcoal (50 g) can be considered if ingested within 2-4 hours • Hemodialysis could be considered if drug level is high, especially in patients with poor renal function. • Fresh-frozen plasma is not recommended for DOAC reversal 	<ul style="list-style-type: none"> • Apix/Riva: Admin ANDEXXA per package insert • Betrix/Edox: Admin off-label ANDEXXA* (800 mg at 30 mg/min then 8 mg/min for up to 120 min)⁴ • Admin 4F-PCC 50 units/kg IV (if ANDEXXA not avail/used) • If 4F-PCC is not available, consider aPCC 50 units/kg IV (refer to prescribing information for max units) • Consider Activated charcoal (50 g) if ingested <24 hrs • Fresh-frozen plasma is not recommended for DOAC reversal

4F-PCC= four-factor prothrombin complex; aPCC= activated prothrombin complex concentrate; *Off-label ANDEXXA OR 4F-PCC suggested for Betrix/Edox⁴

Restart DOAC

- Most patients benefit from restarting anticoagulation after bleeds, but make sure there is still a valid indication.
 - eg. CHA₂DS₂-VASc is ≥ 1 (in AF), length of treatment hasn't been reached (for VTE treatment or post-op prophylaxis).
- Base plan on the balance between bleeding and thromboembolic risks and discussions with other appropriate practitioners (eg. surgeons), the patient, and caregivers.
- Timing of restart: Delay restart if bleeding occurred in a critical site or if patient has a high risk for re-bleeding. Patients with GI bleed should typically wait at least 7-14 days. Patients with intracranial hemorrhage (and no mechanical valve) should wait at least 4 weeks.² In patients with moderate to high risk of recurrent VTE without high risk of recurrent bleeding, ASH suggests resuming anticoagulation within 90 days rather than discontinuation.³
- Make sure dose is correct based on age, renal function, weight, and indication and address any reversible risk factors such as interacting medications or unnecessary antiplatelet therapy.

References

- Unless otherwise referenced, document adapted from: 2017 ACC Expert Consensus Decision Pathway on Management of Bleeding in Patients on Oral Anticoagulants. JACC, Volume 70, Issue 24, 19 December 2017, Pages 3042-3067, ISSN 0735-1097, <https://doi.org/10.1016/j.jacc.2017.09.1085>
- ¹Levy JH, Ageno W, Chan NC, Crowther M, Verhamme P, Weitz J. When and how to use antidotes for the reversal of direct oral anticoagulants: guidance from the SSC of the ISTH. J Thromb Haemost. 2016 Mar;14(3):623-7. doi: 10.1111/jth.13227. Epub 2016 Feb 17.
- ²Hemphill, et al. 2015 AHA/ASA Guidelines for the Management of Spontaneous Intracerebral Hemorrhage. Stroke. 2015;46:000-000. DOI: 10.1161/STR.0000000000000069
- ³Witt WM, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: optimal management of anticoagulation therapy. DOI 10.1182/bloodadvances.2018024893
- ⁴Cuker A, et al. Reversal of direct oral anticoagulants: Guidance from the Anticoagulation Forum. Am J Hematol.2019;94:697–709. doi.org/10.1002/ajh.25475

Disclaimer: This document is for informational purposes only and does not, itself, constitute medical advice. This document is not a replacement for careful medical judgments by qualified medical personnel. There may be information in this document that does not apply to or may be inappropriate for the medical situation at hand.